



TITLE:

On optimal 2-uniform convexity inequalities (Nonlinear Analysis and Convex Analysis)

AUTHOR(S):

Kato, Mikio; Takahashi, Yasuji

CITATION:

Kato, Mikio ...[et al]. On optimal 2-uniform convexity inequalities (Nonlinear Analysis and Convex Analysis). 数理解析研究所講究録 2002, 1298: 65-69

ISSUE DATE:

2002-12

URL:

<http://hdl.handle.net/2433/42683>

RIGHT:

On optimal 2-uniform convexity inequalities

九州工大工 加藤幹雄 (Mikio Kato)

岡山県立大情報工 高橋泰嗣 (Yasuji Takahashi)

This is a résumé of some recent results of the authors on optimal 2-uniform convexity inequalities.

A Banach space X is called q -uniformly convex ($2 \leq q < \infty$) if there is $C > 0$ such that

$$\delta_X(\varepsilon) \geq C\varepsilon^q \text{ for all } \varepsilon > 0, \quad (1)$$

where $\delta_X(\varepsilon)$ is the modulus of convexity,

$$\delta_X(\varepsilon) = \inf \left\{ 1 - \left\| \frac{x+y}{2} \right\| : \|x\| = \|y\| = 1, \|x-y\| = \varepsilon \right\}. \quad (2)$$

The q -uniform convexity of X is characterized by the following " q -uniform convexity inequality"

$$\frac{\|x+y\|^q + \|x-y\|^q}{2} \geq \|x\|^q + \|Cy\|^q, \quad (3)$$

where $0 < C \leq 1$, independent on $x, y \in X$ (cf. [1,2,4]).

Clarkson's inequalities imply that L_q ($2 \leq q < \infty$) is q -uniformly convex and L_p ($1 < p \leq 2$) is p' -uniformly convex, where $1/p + 1/p' = 1$, whereas, as is well known, L_p ($1 < p \leq 2$) is in fact 2-uniformly convex; Ball-Carlen-Lieb [1] gave a proof which uses Hanner's and Gross' inequality. The *optimal 2-uniform convexity inequality* for L_p ($1 < p \leq 2$) is the following:

$$\frac{\|f+g\|_p^2 + \|f-g\|_p^2}{2} \geq \|f\|_p^2 + (p-1)\|g\|_p^2, \quad (4)$$

where the constant $p-1$ is optimal. This is equivalent to the following more sharp inequality

$$\left(\frac{\|f+g\|_p^p + \|f-g\|_p^p}{2} \right)^{1/p} \geq \left(\|f\|_p^2 + (p-1)\|g\|_p^2 \right)^{1/2}, \quad (5)$$

where $p-1$ is optimal ([1]). (For $2 \leq p < \infty$ these inequalities are reversed; see Ball-Carlen-Lieb [1].) The inequality (5) yields the following best estimate in (1) for L_p ($1 < p \leq 2$):

$$\delta_{L_p}(\varepsilon) \geq \{(p-1)/8\}\varepsilon^q \text{ for all } \varepsilon > 0.$$

In the recent paper [5] Takahashi-Hashimoto-Kato presented some generalizations of the q -uniform convexity inequality (3), and showed that these inequalities are inherited to the Lebesgue-Bochner space $L_r(X)$. In this note, by using their results, we shall present some generalizations of the optimal 2-uniform convexity inequalities (4) and (5).

First we state the following inequalities which are fundamental in our discussion:

Lemma 1 ([4, p.76]). Let $1 < p \leq q < \infty$ and $\gamma = \sqrt{(p-1)/(q-1)}$. Then:

(i) For any $x, y \in X$

$$\left(\frac{\|x+y\|^p + \|x-y\|^p}{2} \right)^{1/p} \leq \left(\frac{\|x+y\|^q + \|x-y\|^q}{2} \right)^{1/q} \quad (6)$$

(ii) For any $x, y \in X$

$$\left(\frac{\|x+\gamma y\|^q + \|x-\gamma y\|^q}{2} \right)^{1/q} \leq \left(\frac{\|x+y\|^p + \|x-y\|^p}{2} \right)^{1/p} \quad (7)$$

Theorem 1 (Takahashi-Hashimoto-Kato [5]). Let $2 \leq q < \infty$ and $1 < t \leq \infty$. The following are equivalent.

(i) X is q -uniformly convex.

(ii) For any $1 < t \leq \infty$ there exists $0 < C \leq 1$ such that

$$\left(\frac{\|x+y\|^t + \|x-y\|^t}{2} \right)^{1/t} \geq (\|x\|^q + \|Cy\|^q)^{1/q} \quad \forall x, y \in X. \quad (8)$$

(iii) For some $1 < t \leq \infty$ there exists $0 < C \leq 1$ such that the inequality (8) holds.

In particular, we have

Theorem 2 (2-uniform convexity inequalities). The following are equivalent.

(i) X is 2-uniformly convex.

(ii) For any $1 < t \leq \infty$ there exists $0 < C \leq 1$ such that

$$\left(\frac{\|x+y\|^t + \|x-y\|^t}{2} \right)^{1/t} \geq (\|x\|^2 + \|Cy\|^2)^{1/2} \quad \forall x, y \in X. \quad (9)$$

(iii) For some $1 < t \leq \infty$ there exists $0 < C \leq 1$ such that (9) holds.

Remark 1. In Theorem 2 (ii) and (iii) we have $0 < C \leq \min\{1, t-1\}$, where equality holds if X is a Hilbert space.

Proposition 1. Assume that the following 2-uniform convexity inequality

$$\max\{\|x + y\|, \|x - y\|\} \geq (\|x\|^2 + C\|y\|^2)^{1/2} \quad (10)$$

holds in X . Then,

$$\delta_X(\epsilon) \geq \frac{C}{8}\epsilon^2 \quad \text{for all } 0 < \epsilon < 2. \quad (11)$$

One should note that for $1 < t < \infty$

$$\max\{\|x + y\|, \|x - y\|\} \geq \left(\frac{\|x + y\|^t + \|x - y\|^t}{2} \right)^{1/t}.$$

Now, 2-uniform convexity inequality is inherited to $L_r(X)$ as follows.

Theorem 3. Let $1 < p, r \leq 2$. Assume that the inequality

$$\left(\frac{\|x + y\|^p + \|x - y\|^p}{2} \right)^{1/p} \geq (\|x\|^2 + C\|y\|^2)^{1/2} \quad (12)$$

holds in X . Then

$$\left(\frac{\|f + g\|_r^p + \|f - g\|_r^p}{2} \right)^{1/p} \geq (\|f\|_r^2 + C'\|g\|_r^2)^{1/2} \quad (13)$$

holds in $L_r(X)$, where

$$C' = \begin{cases} C & \text{if } p \leq r \leq 2, \\ \{(r-1)/(p-1)\}C & \text{if } 1 < r < p. \end{cases}$$

Remark 2. The constant C' is optimal in general.

Since X is isometrically embedded into $L_r(X)$, it is trivial that any inequality valid in $L_r(X)$ holds in X . The next result asserts that from a 2-uniform convexity inequality in $L_r(X)$ we have a stronger one in X .

Theorem 4. Let $1 < r \leq 2$ and $r < p$. Assume that

$$\left(\frac{\|f + g\|_r^p + \|f - g\|_r^p}{2} \right)^{1/p} \geq (\|f\|_r^2 + C\|g\|_r^2)^{1/2} \quad (14)$$

holds in $L_r(X)$. Then

$$\left(\frac{\|x + y\|^r + \|x - y\|^r}{2} \right)^{1/r} \geq (\|x\|^2 + C\|y\|^2)^{1/2} \quad (15)$$

holds in X .

Indeed take any non-zero $x, y \in X$ and put $f = (x, x)$, $g = (y, -y) \in \ell_r^2(X) \subset L_r(X)$ in (14).

By Theorems 3 and 4 we have the following optimal 2-uniform convexity inequality for L_r (use the parallelogram law for scalars).

Theorem 5 (Optimal 2-uniform convexity inequality for L_r , $1 < r \leq 2$). Let $1 \leq r \leq 2$ and $1 < p \leq \infty$. Then

$$\left(\frac{\|f + g\|_r^p + \|f - g\|_r^p}{2} \right)^{1/p} \geq \left(\|f\|_r^2 + C\|g\|_r^2 \right)^{1/2} \quad (16)$$

holds in L_r , where $C = \min\{p - 1, r - 1\}$.

Remark 3. (i) The constant C in (16) is best possible.

(ii) The inequality (16) for L_p , $1 < p \leq 2$ with $C = p - 1$, that is,

$$\left(\frac{\|f + g\|_p^p + \|f - g\|_p^p}{2} \right)^{1/p} \geq \left(\|f\|_p^2 + (p - 1)\|g\|_p^2 \right)^{1/2} \quad (5)$$

was proved in Ball-Carlen-Lieb [1]. Their proof used Hanner's inequality and Gross' inequality, whereas we derived (5) from Theorems 3 and 4 and the parallelogram law for scalars.

Theorem 3 yields the following

Theorem 6 (Optimal 2-uniform convexity inequality for $L_r(L_s)$, $1 < r, s \leq 2$). Let $1 \leq r, s \leq 2$ and $1 < p \leq \infty$. Then

$$\left(\frac{\|f + g\|_r^p + \|f - g\|_r^p}{2} \right)^{1/p} \geq \left(\|f\|_r^2 + C\|g\|_r^2 \right)^{1/2} \quad (17)$$

holds in $L_r(L_s)$, where $C = \min\{p - 1, r - 1, s - 1\}$. In particular, if $1 < p \leq \min\{r, s\}$, then $C = p - 1$.

Remark 4. The constant C in (17) is best possible.

References

- [1] K. Ball, E. A. Carlen and E. H. Lieb, Sharp uniform convexity and smoothness inequalities for trace norms, *Invent. Math.* **115** (1994), 463–482.
- [2] B. Beauzamy, *Introduction to Banach Spaces and Their Geometry* 2nd Ed., 1985.

- [3] M. Kato and Y. Takahashi, Type, cotype constants and Clarkson's inequalities for Banach spaces, *Math. Nachr.* **186** (1997), 187–196.
- [4] J. Lindenstrauss and L. Tzafriri, *Classical Banach Spaces II*, 1979.
- [5] Y. Takahashi, K. Hashimoto and M. Kato, On sharp uniform convexity, smoothness, and strong type, cotype inequalities, *J. Nonlinear Convex Anal.* **3** (2002), 267–281.
- [6] Y. Takahashi and M. Kato, Clarkson and Random Clarkson inequalities for $L_r(X)$, *Math. Nachr.* **188** (1997), 341–348.

*Department of Mathematics,
Kyushu Institute of Technology,
Kitakyushu 804-8550, Japan
e-mail: katom@tobata.isc.kyutech.ac.jp*

*Department of System Engineering,
Okayama Prefectural University,
Soja 719-1197, Japan
e-mail: takahasi@cse.oka-pu.ac.jp*